

Title: Intrinsic connectivity in the human brain does not reveal networks for “basic” emotions

Running title: Intrinsic connectivity of emotion

Alexandra Touroutoglou^{1,2,3}, Kristen A. Lindquist^{4,5}, Bradford C. Dickerson^{2,6}, Lisa Feldman Barrett^{2,3,7*}

¹ Department of Neurology, Massachusetts General Hospital and Harvard Medical School, Charlestown, MA, USA

²Athinoula A. Martinos Center for Biomedical Imaging, Massachusetts General Hospital and Harvard Medical School, Charlestown, MA, USA

³Psychiatric Neuroimaging Division, Department of Psychiatry, Massachusetts General Hospital and Harvard Medical School, Charlestown, MA ,USA

⁴Department of Psychology, University of North Carolina, Chapel Hill, NC, USA

⁵ Biomedical Research Imaging Center, University of North Carolina, Chapel Hill, NC, USA

⁶Frontotemporal Disorders Unit, Department of Neurology, Massachusetts General Hospital and Harvard Medical School, Charlestown, MA, USA

⁷Department of Psychology, Northeastern University, Boston, MA, USA

* Correspondence: Dr. Lisa Feldman Barrett, Department of Psychology, 125 Nightingale Hall, Northeastern University, Boston, MA 02115-5000, USA

Abstract

We tested two competing models for the brain basis of emotion, the basic emotion theory and the conceptual act theory of emotion, using resting-state functional connectivity magnetic resonance imaging (rs-fcMRI). The basic emotion view hypothesizes that anger, sadness, fear, disgust and happiness each arise from a brain network that is innate, anatomically constrained, and homologous in other animals. The conceptual act theory of emotion hypothesizes that an instance of emotion is a brain state constructed from the interaction of domain-general, core systems within the brain such as the salience, default mode, and frontoparietal control networks. Using peak coordinates derived from a meta-analysis of task-related emotion fMRI studies, we generated a set of whole-brain rs-fcMRI “discovery” maps for each emotion category, and examined the spatial overlap in their conjunctions. Instead of discovering a specific network for each emotion category, variance in the discovery maps was accounted for by the known domain-general network. Furthermore, the salience network observed as part of every emotion category. These results indicate that specific networks for each emotion do not exist within the intrinsic architecture of the human brain, and instead support the conceptual act theory of emotion.

Keywords: emotions, intrinsic functional connectivity networks, basic emotion theory, conceptual act theory of emotion

1. Introduction

For the past fifty years, scientists have been largely convinced that certain emotions, such as anger, fear, sadness, happiness and disgust, are biologically basic, meaning that they are natural kinds (Barrett, 2006). In this view, each emotion category arises from an innate, specific brain module with homology to other animals [e.g., (Ekman, 1999; Keltner and Ekman, 2000; Tracy and Randles, 2011)]ⁱ. This basic emotion view has dominated the science of emotion, and is widely accepted in the popular media, despite the fact the brain basis of emotion is still poorly understood. Although careful and elegant studies of so-called “emotional” behavioral adaptations in non-human animals have revealed distinct neural circuits that control escape(Vazdarjanova and McGaugh, 1998), freezing (LeDoux, 2007), fighting [e.g., offensive attack (Lin, et al., 2011); defensive aggression (Motta, et al., 2009)] there are a number arguments for why a neural circuit for a behavior cannot be considered a neural circuit for an emotion per se [e.g., (Barrett, 2012; Barrett, et al., 2007; LeDoux, 2012)]. For example, depending on the circumstances, an animal might flee, freeze, or fight when faced with potential danger (i.e., during a “fearful” situation). This introduces the problem of having many fear circuits [e.g., (Gross and Canteras, 2012)] and poses an inductive problem for the science of basic emotions. Cognitive neuroscience and lesion research has searched for emotion brain modules with little success [for recent meta-analytic evidence, see (Kober, et al., 2008; Lindquist, et al., 2012); for a discussion, see (Barrett and Satpute, 2013; Lindquist and Barrett, 2012)]. One recent meta-analysis of the human neuroimaging literature was interpreted as supportive of the basic emotion hypothesis (Vytal and Hamann, 2010), but in fact reported limbic and non-limbic regions as showing consistent but non-specific increases in activation during anger, sadness, fear, disgust and happiness [for an alternative interpretation of their findings see (Hamann, 2012; Lindquist, et

al., 2012)].

In contrast to the basic emotion view, a constructionist approach to emotion, the “conceptual act theory of emotion” hypothesizes that an emotion such as anger, sadness, fear, disgust, or happiness is a population of instances; the instances do not arise from their own, dedicated brain network, but are instead *constructed* from the combination of activity in domain-general, core brain systems that perform more basic psychological functions such as salience detection, memory, sensory perception, language, and so on [(Barrett, 2006; Barrett, 2012; Barrett and Satpute, 2013; Lindquist and Barrett, 2012)]. In particular, the conceptual act theory of emotion predicts that the same intrinsic networks would be engaged during a variety of emotions, although perhaps in different patterns [for an extension of this view see (Oosterwijk, et al., 2012)]. Indeed, preliminary support for the conceptual act theory of emotion view can be observed in several recent neuroimaging experiments (Wilson-Mendenhall, et al., 2013; Wilson-Mendenhall, et al., 2011) demonstrating the involvement of brain areas associated with representing body states, salient events, memory, sensory perception, and language during emotional experiences. Other evidence consistent with the conceptual act theory of emotion comes from intracranial stimulation studies (Guillory and Bujarski, 2014), as well as meta-analyses of neuroimaging experiments demonstrating that domain-general brain systems are commonly engaged across a variety of emotion categories (Lindquist, et al., 2012) and during both emotional and cognitive events (Barrett and Satpute, 2013; Lindquist and Barrett, 2012).

In this study, we compared the “basic emotion” and “conceptual act” theories using an analysis of the brain’s intrinsic functional connectivity. A recent explosion of research demonstrates that the human brain contains a small world architecture with densely connected “hubs” (Sporns, 2013; van den Heuvel and Sporns, 2013). Within this structure, the brain shows

continuous, intrinsic activity organized as connected networks. These networks, referred to as “intrinsic networks,” are identified in temporal correlations of the low frequency blood oxygen level-dependent (BOLD) signal fluctuations in voxels while a participant lays “at rest” during functional magnetic resonance imaging (called “resting state functional connectivity magnetic resonance imaging” or rs-fcMRI). Critically, patterns of intrinsic activity are quite similar to the patterns of task-related activity [(Smith, et al., 2009; Spector, et al., 2009); for a review see (Bressler and Menon, 2010)]. These networks also account for a large proportion of the brain’s metabolic budget (Raichle, 2010). Together, these findings have thus led researchers to conclude that spontaneous neuronal activity within these networks reflects the intrinsic organization of the brain (Buckner, et al., 2013), which in turn forms the functional architecture of the mind (Barrett and Satpute, 2013; Lindquist and Barrett, 2012; Menon, 2011).

There are several reasons why intrinsic brain networks are useful for comparing the basic emotion and conceptual act theories. First, intrinsic brain networks have anatomic and cross-species properties that make rs-fcMRI an ideal approach for testing the basic emotion theory’s hypothesis that dedicated emotion networks exist. Intrinsic networks are anatomically constrained (Deco, et al., 2010; Hermundstad, et al., 2013; Pernice, et al., 2011; van den Heuvel, et al., 2009), and can be observed under anesthesia (Greicius, et al., 2008). The networks are found in people of different cultures [e.g., (Manoliu, et al., 2013; Wang, et al., 2014)]. Several of the networks are homologous with the networks that exist within the brains of other animals (Hayes and Northoff, 2011; Mantini, et al., 2013; Rilling, et al., 2007; Vincent, et al., 2007). By comparison, basic emotion theory hypothesizes that (1) each emotion is caused by a specific, dedicated network that is anatomically intrinsic to the human brain, (2) each network should be universal, and (3) homologous in non human animals. Furthermore, some theories propose that

these networks should be confined to subcortical regions of the brain [e.g. (Panksepp, 1998)], making it difficult to identify them in studies of humans where direct anatomical investigations are difficult. Second, many rs-fcMRI studies reveal the existence of domain-general intrinsic networks; these can be used to test the conceptual act theory's hypothesis that an emotion is constructed as an interaction of domain-general systems. To date, networks have been identified for salience (Seeley et al., 2007; Touroutoglou et al., 2012), language (Lohmann, et al., 2010; Tomasi and Volkow, 2012), executive function (Seeley et al., 2007; Vincent et al., 2008), attention (Corbetta et al., 2008; Vincent et al., 2008), semantic processing (Binder, et al., 2009), memory (Buckner, et al., 2008; Dickerson and Eichenbaum, 2010) and other processes [(Yeo, et al., 2011) for a review see (Bressler and Menon, 2010)]. If emotions are constructed as interactions of basic networks that subserve domain-general functions in the brain, then rs-fcMRI analyses should reveal (1) evidence of many intrinsic networks contributing to a single instance of an emotion category and (2) evidence of a common intrinsic network contributing to different emotion categories. Thus, the rs-fcMRI approach, while not a perfect window into the structure within the human brain, does allow an opportunity to investigate the structural properties of the brain in relation to emotion.

Although a number of intrinsic networks have now been replicated across samples and analysis methods [e.g., seed-based rs-fcMRI (Vincent, et al., 2008); independent components analysis (Smith, et al., 2009)], no studies to date have explicitly used rs-fcMRI to examine whether a specific intrinsic connectivity network can be identified for each emotion category.

In the present study, we thus used a “seed and discover” method (Vincent, et al., 2008) to assess whether there are specific intrinsic networks for specific emotions or whether the networks underlying emotional experiences and perceptions are comprised as combinations of

domain-general intrinsic networks such as those found in Shirer et al. (2012), Yeo et al. (2011) and Smith et al. (2009) (e.g., networks involved in salience detection, memory, attention, language, motor function and sensation). The “seed and discover method” relied on two types of evidence: meta-analytic data and resting-state data. (1) Meta-analytic peaks of task-evoked fMRI activity; these identified a priori regions of interest (ROIs) that consistently showed an increase in activity during emotional experiences and emotion perception. (2) These peaks were then used as “seeds” in the analysis of rs-fcMRI data from two samples to generate the intrinsic “discovery” maps for each emotion category (Sample 1, N = 89; Sample 2, N = 300). In particular, we extracted the time course of BOLD activity from a predefined “seed” region and compared it to the timecourse of all other voxels in the brain. The result was a “discovery” map of voxels that showed a similar BOLD response across time. Voxels whose timecourses correlate significantly with one another are considered to be part of the same rs-fcMRI intrinsic network.

To investigate whether intrinsic networks exist for anger, disgust, fear, sadness, and happiness, we chose ROI seeds from the activation peaks reported in Vytal and Hamann’s (2010) meta-analysis of task-related functional neuroimaging studies of emotion experience and perception (see Table 1). In contrast to resting state functional connectivity analyses, task-evoked fMRI studies reveal brain regions that show increases in activation (relative to some baseline condition) during specific psychological tasks or conditions (e.g., experiencing anger relative to a neutral emotional state). Meta-analyses of task-evoked activity overcome the Type I error prevalent in individual task-evoked fMRI studies by revealing those brain areas that consistently show increases in activity during a particular condition (e.g., anger) across studies (cf. Lindquist et al. 2012). Using the activation likelihood estimation (ALE) meta-analytic method, Vytal and Hamann (2010) thus identified regions that were consistently activated across studies for each of

five emotion categories (anger, sadness, fear, disgust, and happiness).

If anatomically-constrained networks for each emotion category exist in the intrinsic architecture of the human brain, as the basic emotion view predicts, then the meta-analytically derived seed regions for a given emotion category (i.e., the peaks of consistent activation for a given category of emotion, such as happiness) should produce “discovery” maps whose spatial overlap reveals a network for that category. This finding would provide strong support for the hypothesis that emotions are biologically basic categories reflected in the intrinsic structure of the brain. Alternatively, if the peaks observed in Vytal and Hamann (2010) are nodes in domain-general intrinsic networks, as predicted by the conceptual act theory of emotion, then the conjunction of the discovery maps for a given emotion category would not converge on a single network. Instead, emotion-based seeds would give evidence of the domain-general intrinsic networks that are already known to exist in the literature.

To ascertain the degree of spatial overlap between the discovery maps that were generated by the meta-analytically derived emotion seeds and the maps for well-known domain-general intrinsic networks (i.e., visual, language, episodic memory, executive function, salience detection networks) (Shirer, et al., 2012; Yeo, et al., 2011), we used a goodness-of-fit metric (Greicius, et al., 2004). The conceptual act theory of emotion hypothesizes that emotional instances are constructed from the interactions of these domain-general networks. According to this hypothesis, the brain predicts incoming sensory input from the body and the world by categorizing it, thereby constructing it into meaningful emotional experiences and perceptions (Barrett, 2009; Barrett, 2012; Barrett, 2013; Barrett, 2014; Lindquist and Barrett, 2012). From this perspective, the salience, default mode network, and frontoparietal control networks are centrally important to constructing instances emotion (being involved in interoception, semantic

processing, and categorization, respectively), but so too are the exteroceptive and other attention networks as well. An instance of emotion is understood as a brain state, constructed as the ongoing interaction of brain networks.

Finally, we also assessed the conceptual act theory of emotion hypothesis that all emotions constructed with the “salience network.” The strength of connectivity within this network is correlated to the intensity of affective experience (Seeley, et al., 2007; Touroutoglou, et al., 2014; Touroutoglou, et al., 2012) and nodes within this network show an increase in activation across varying instances of emotions in task-evoked studies (Wilson-Mendenhall, et al., 2013). Furthermore, this network is particularly relevant to negative emotions (Bickart, et al., 2012; Hayes and Northoff, 2011; Seeley, et al., 2007; Touroutoglou, et al., 2014; Touroutoglou, et al., 2012). To test this hypothesis, we used the same seed and discover method to determine whether there was any spatial overlap between the “discovery” maps for anger, sadness, fear, disgust. Support for basic emotion theory would be found if the different negative emotion categories each had a distinct network. Support for the conceptual act theory of emotion would be found if the different negative emotion categories had overlapping regions within a common network such as the salience network.

2. Materials and methods

2.1. Participants

Sample 1 consisted of 89 young adults (44 men) ranging in age from 18 to 33, with a mean age of 22.4 years ($SD = 3.34$). Sample 2 consisted of 300 young adults (150 men) with a mean age of 22.3 years ($SD = 1.94$) (rs-fcMRI data from both samples have been previously published in Yeo et al., 2011). All participants were right-handed, native English speakers and had normal or

corrected-to-normal vision. No participant reported a history of neurological or psychiatric disorder.

2.1.1. fMRI data acquisition and preprocessing procedures

Data were collected with a 3 Tesla Tim Trio System (Siemens Medical Systems, Erlangen, Germany), using a 12-channel phased-array head coil. Structural data in Sample 1 and 2 were acquired using a 3D T1-weighted magnetization-prepared gradient-echo image (MPRAGE) [repetition time (TR) = 2200 ms; echo time (TE) = 1.54 ms; flip angle (FA) = 70°, 1.2 mm isotropic voxels]. Whole-brain fMRI data were acquired with echo-planar sequence [Sample 1 and Sample 2: TR = 3000 ms; TE = 30 ms; FA = 90°; 3.0 mm isotropic voxels, 47 slices]. During the resting-state fMRI runs, participants were instructed to keep their eyes open. Head motion was minimized using head restraints, including a pillow and foam padding. Noise was attenuated with ear plugs.

Preprocessing of the fMRI data involved a series of previously established rs-fcMRI procedures (Van Dijk, et al., 2010), including: (1) removal of the first four volumes to allow for T1 equilibration effects, (2) slice timing correction (SPM2, Wellcome Department of Cognitive Neurology, London, United Kingdom), and (3) head motion correction (FMRIB, Oxford, UK). Data were normalized to the Montreal Neurological Institute (MNI) atlas space (SPM2, Wellcome Department of Cognitive Neurology, London, United Kingdom) and re-sampled to 2mm cubic voxels. A low-pass temporal filter removed frequencies higher than 0.08Hz. Data were spatially smoothed using a 6 mm full-width half-maximum Gaussian filter. Sources of spurious variance and their temporal derivatives were removed through linear regression

including: (1) six parameters obtained by rigid-body correction of head motion correction, (2) the signal averaged over the whole brain, (3) the signal averaged over the ventricles, and (4) the signal averaged over the deep cerebral white matter.

2.2. Functional connectivity analysis

2.2.1. Selection of Regions of Interest (ROIs)

The atlas coordinates of all seed ROIs are presented in Table 1. To examine whether emotion seed ROIs reveal intrinsic emotion networks, we used peak activations that were more consistent than expected by chance for each emotion within the Vytal and Hamann (2010) meta-analysis (for the validity of this “seed and discover” method see Supplementary Materials). As the seeds for the discovery maps, we selected the peak activations of three different regions with the largest cluster of activation for each emotion category. Our rationale was that these ROIs would constitute the most spatially discriminable regions for each presumed “basic” emotion. For ease of reporting, we only present the analysis using the three largest peak activations (i.e., those showing the highest degree of consistency among individual studies in the literature) reported for each emotion in Vytal and Hamann (2010). To ensure that the number of peaks analyzed did not affect the analysis, we also performed a “seed and discover” analysis using all peak activations reported for each emotion in Vytal and Hamann (2010) (9-19 peaks per emotion) (see Supplementary Fig. 2, 3, and 4).

[INSERT TABLE 1 ABOUT HERE]

2.2.2. Overlapping spatial topography of the rs-fcMRI discovery maps

To create each “discovery” map, we created spherical ROIs (4-mm radius) around each seed region and then computed Pearson’s product moment correlations (r) between the mean signal timecourse of each seed region and the time course of all other voxels in the brain. The resulting correlation maps were converted to z -values, using Fisher’s r -to- z transformation and were averaged across participants. To explore whether each set of seed regions revealed a unified intrinsic connectivity network, we next computed a spatial conjunction analysis on the discovery maps for each emotion category. Specifically, the group-level z -score maps for the seed reference regions were binarized at a minimum threshold of $z(r) = 0.25$. We then computed their convergent spatial overlap, identifying voxels with $z(r)$ values ≥ 0.25 in all seed regions. All maps are shown on slices in MNI atlas space using the FSL view toolbox (<http://www.fmrib.ox.ac.uk/fsl>) (for the validity of this method see Supplementary Materials).

In addition to generating spatial convergence maps, we quantified the strength of functional connectivity between the seed regions by calculating Fisher’s r -to- z correlation coefficients between each pair of seeds. Next, we calculated the average connectivity measure of $z(r)$ values between the seeds associated with the three seeds for each emotion category. As a reference range for functional connectivity strength values, we calculated the average connectivity measure of $z(r)$ values within the well-known “default mode network” (DMN) (e.g., Andrews-Hanna, et al., 2010) (see Supplementary Materials). As a control analysis, we selected seed ROIs in visual, motor and auditory cortex that are typically uncorrelated (as in Van Dijk et al., 2010) (see Supplementary Materials).

To test the hypothesis that seed regions from each negative emotion were part of the same intrinsic rs-fcMRI network [i.e., the salience network (Seeley, et al., 2007)], we examined

the conjunction of the discovery maps for the single largest (i.e., most spatially distinctive) meta-analytic peak for each discrete negative emotion (see Table 1).

2.2.3. Goodness-of-fit analysis between the rs-fcMRI emotion maps and primary intrinsic connectivity networks

To test the hypothesis that the emotion seed regions produced rs-fcMRI maps that were representative of canonical intrinsic connectivity networks, we calculated a goodness-of-fit metric (Greicius, et al., 2004) that represented a spatial similarity index over the entire map. For this analysis, we chose a set of 14 intrinsic connectivity networks identified by Shirer et al. (2012) (Table 2; see also Figure 2S in Shirer et al (2012). Because the “salience network” plays a critical role in affective experience (Seeley, et al., 2007; Touroutoglou, et al., 2014; Touroutoglou, et al., 2012), we included the dorsal salience subnetwork (most clearly involving connections between the dorsal anterior insula and dorsal ACC) and the ventral salience subnetwork network (involving connections between ventral anterior insula and pregenual ACC extending to the subgenual ACC) identified by Touroutoglou et al. (2012). The canonical intrinsic connectivity networks of interest included in this analysis are presented in Table 2.

A template of each of the intrinsic connectivity networks of interest was used to select the “best-fit” of rs-fcMRI emotion maps. We used the template-matching procedure developed by Greicius et al. (2004) that involved taking the average z score of voxels falling within the template minus the average z score of voxels outside the template and selecting the network of interest in which this difference (the goodness-of-fit) was the greatest. As a reference range for goodness-of-fit values, we calculated the goodness-of-fit metric between the rs-fcMRI DMN maps and the canonical intrinsic connectivity networks (see Supplementary Materials). We

expected the rs-fcMRI DMN discovery maps to have high fit values with dorsal and ventral DMN but low fit values with the sensorimotor, auditory, or visuospatial networks identified by Shirer et al. (2012).

[INSERT TABLE 2 ABOUT HERE]

2.2.4. Reliability of rs-fcMRI emotion maps

To assess the reliability of the strength of connectivity of the rs-fcMRI emotion maps, we computed intraclass correlation coefficients (ICC) (two way random effects with absolute agreement) between the connectivity $z(r)$ values between the trio of seeds associated with each emotion category in Sample 1 and in Sample 2, using PASW Statistics 18, Release Version 18.0.0 (SPSS, Inc., 2009, Chicago, IL, www.spss.com).

3. Results

3.1. Peaks of consistent activity during emotion belong to domain-general intrinsic networks

Inconsistent with the basic emotion hypothesis and consistent with the conceptual act theory of emotion hypothesis, we did not find strong evidence for intrinsic networks corresponding to specific emotions (Fig. 1). For instance, the anger peak discovery maps (created from three of the largest meta-analytic activation peaks from Vytal and Hamann, 2010 and the voxels correlated with each peak) did not share spatial overlap with one another. As a result, the conjunction of the anger discovery maps was empty, indicating that an anatomically-constrained network for anger does not exist within the intrinsic architecture of the human brain. Neither “discovery” maps at our a priori threshold of $z(r) = 0.25$, nor maps at a less stringent threshold of $z(r) = 0.1$, revealed

an intrinsic network. We repeated this analysis, with the same result, for the peaks that consistently activated in Vytal and Hamann's analysis during sadness, fear, disgust, and happiness. The average connectivity strength, $z(r)$, between the seed regions for the five emotions are shown in Figure 2. Furthermore, when we repeated the analysis using a larger number of seeds from all the meta-analytic peaks available in Vytal and Hamann (2010), the results did not change. Specifically, thirteen peaks that were consistently activated during anger, nineteen peaks that were consistently activated during sadness, ten peaks that were consistently activated during fear, sixteen peaks that were consistently activated for disgust and nine peaks that were consistently activated during happiness in Vytal and Hamann's (2010) meta-analysis did not together reveal an intrinsic network for each emotion category (see Supplementary Fig. 2 and 3).

Instead, the goodness-of-fit analysis revealed that Vytal and Hamann (2010)'s peaks were nodes in the set of domain-general intrinsic networks already identified in the literature (Table 3; see also Supplementary Fig. 4). Within each emotion category, the discovery maps did not show convergent overlap with just a single intrinsic network. For example, the superior temporal gyrus ROI that showed consistent increases in activity across studies of happiness was part of the visuospatial intrinsic network and the high-level visual network. By contrast, the left pregenual anterior cingulate cortex ROI that showed consistent increases across studies of happiness was part of the ventral salience network and dorsal DMN (see Table 3). Critically, intrinsic networks were identified in the discovery maps of multiple negative emotions categories. As an example, the domain-general dorsal salience network was identified in the discovery maps for different negative emotion categories, i.e., fear, disgust, and sadness. The dorsal extent of the default

mode network was also identified in the discovery maps for all emotion categories (see Supplementary Fig. 4).

Finally, as predicted by the conceptual act theory of emotion hypothesis, our seed and discovery method for the negative emotion categories revealed the salience network. Spatial overlap between discovery maps derived from the largest meta-analytic peak for each negative emotion revealed regions within the salience network, such ventral anterior insula, caudate, and thalamus (Fig. 3). To further examine whether the ventral anterior insula, caudate, and thalamus were indeed evidence of the broader salience network, we performed an exploratory analysis where we lowered the threshold to $z(r) = 0.05$. At this lower threshold, it became clear that the activity in ventral anterior insula, caudate, and thalamus that we observed was indeed part of the canonical salience network (Seeley et al., 2007; Touroutoglou et al., 2012). Of note, we did not formally address whether seeds from happiness converged on the salience network, as there was only one positive emotion in our analysis. Nonetheless, as seen in the goodness of fit analysis (see Table 3), the happiness intrinsic connectivity map (anchored by the ACC) included regions that also overlapped with the ventral salience network.

[INSERT FIGURES 1, 2, 3 AND TABLE 3 ABOUT HERE]

3.2. Reliability of rs-fcMRI emotion maps

All networks showed high reliability across Samples 1 and 2, supporting the generalizability of our observations that basic emotion networks do not exist within the intrinsic architecture of the human brain. Most importantly, the intraclass correlation coefficients (ICC) across the two samples for the connectivity $z(r)$ values between the trio of seeds associated with each emotion

demonstrated high reliability for happiness (ICC = 0.98, two-way random effects, $p < 0.02$), fear (ICC = 0.98, two-way random effects, $p < 0.02$), sadness (ICC = 0.99, two-way random effects, $p < 0.09$) disgust (ICC = 0.99, two-way random effects, $p < 0.05$), although they were lower for anger (ICC = 0.65, two-way random effects, $p = \text{ns}$).

4. Discussion

We used an intrinsic connectivity approach to compare two competing hypotheses about the brain basis of emotion. One view hypothesizes that certain emotions (anger, disgust, fear, happiness, and sadness) are biologically “basic” and arise from innate, anatomically constrained brain networks that are homologous in human and non-human animals. Support for this hypothesis would have arisen if we observed anatomically constrained intrinsic networks for specific emotions. This would have been observed if brain regions with consistent increases in activity during emotion experience and perception were each associated with a single intrinsic brain network (e.g., areas that had increased activity during experiences and perceptions of anger were part of an intrinsic network for anger). The alternative theoretical approach, the “conceptual act theory” of emotion, hypothesizes that emotions are constructed from the interaction of domain-general, core systems within the brain. Support for the constructionist hypothesis would arise if we discovered that the regions consistently active during emotional experiences and perceptions were parts of domain-general networks that perform more basic psychological functions. The Vytal and Hamann (2010) meta-analysis identified peak activation coordinates that consistently showed an increase in activation during the experience or perception of a given emotion spanning many studies using many different methods; we asked “do the voxels in these peaks belong to the same intrinsic network in brain, or do they belong to different intrinsic

networks working together?” Furthermore, the seed-based method employed here is sensitive enough to show spontaneous activity of subcortical regions (Bickart, et al., 2012). We therefore had the power to reveal evidence for anatomically-based subcortical networks for each emotion that have been proposed by some basic emotion theorist (e.g., Panksepp, 1998), if they exist.

Using a “seed and discovery” method, we did not find evidence for emotion-specific networks within the intrinsic functional architecture of the brain. Instead, our emotion discovery maps reflected combinations of domain-general networks, such as the salience network, DMN, basal ganglia network, and executive control network, consistent with the hypothesis that different emotions arise from the interaction of domain-general systems within the brain (Barrett, 2012; Barrett and Satpute, 2013; Lindquist and Barrett, 2012). Further evidence for a domain-general constructionist account about emotion comes from our finding that a conjunction of discovery maps for anger, sadness, fear, and disgust each revealed major nodes of the salience network (Seeley et al., 2007) consisting of the anterior insula, caudate, thalamus, and ACC. Consistent with this interpretation, the salience network shows task-evoked activity during the experience of unpleasant affect (Hayes and Northoff, 2011; Lindquist, et al., in press).

Furthermore, individuals with stronger intrinsic connectivity in the salience network report more intense anxiety (Seeley et al., 2007) and arousal when viewing negative images (Touroutoglou et al., 2012). Our findings are also consistent with Laird et al. (2011)’s results showing that a limbic intrinsic connectivity system comprising mostly of limbic and medial temporal regions of the large-scale distributed salience network was associated with the perception of different emotions, i.e., happiness and fear. Together, these findings suggest that the salience network might be playing a general function across instances of anger, disgust, fear, and sadness by representing the feeling of arousal that is common to each of the four emotion categories.

Using this interpretive framework, our results are consistent with other meta-analytic findings (Lindquist et al., 2012) showing that many of the brain regions with consistent increases in activation across studies of *the same* emotion category are, in fact, nodes from different intrinsic networks that have been associated with other basic psychological functions, such as attention, language, memory, salience detection and motor control. Moreover, brain regions with consistent increases during emotion experience and perception can be decomposed into a set of functional groups (i.e., regions that co-activate across studies) (Kober et al., 2008) that resemble the intrinsic networks we observed in the present report. One possibility then, is that intrinsic networks support general psychological processes that form fundamental “ingredients” that contribute to the construction of all manner of mental states (Barrett & Satpute, 2013; Lindquist & Barrett, 2012). For instance, the salience network appears to play a general function across instances of anger, disgust, fear, and sadness (Lindquist et al., 2012) as well as other “emotional” events including empathy (Decety and Jackson, 2004) and autonomic regulation (Craig, 2002; Vogt, 2005). Nodes within the “salience” network are also engaged during “cognitive” events, such as language and executive function tasks [i.e., dorsal anterior insula and dorsal ACC; (Nelson, et al., 2010; Touroutoglou, et al., 2012)] and attention allocation tasks [i.e., dorsal anterior insula and dorsal ACC; (Corbetta, et al., 2008)]. This lack of domain-specificity has led to the interpretation that the salience network functions to orient the brain’s processing capacity towards the most homeostatically-relevant information (constituting a body-based source of attention within the human brain; Lindquist and Barrett, 2012; Barrett and Satpute, 2013) to guide the brain’s “switching” or “reorienting” between “internal” and “external” events (Corbetta, et al., 2008; Menon and Uddin, 2010).

Similarly, nodes within the DMN are engaged during emotion (Lindquist et al., 2012), and also appear to serve more domain-general functions. The DMN nodes are engaged in remembering personal events (autobiographical memory), imagining the future (prospection), accessing memory for word meanings (semantic memory), scene construction and context-based object perception (Binder, et al., 2009) as well as moral reasoning (Bzdok, et al., 2012), and person perception, leading to the suggestion that the DMN creates “situated conceptualizations” (Barrett, 2012; Lindquist et al. 2012; Lindquist & Barrett, 2012) or “mental models” (Barrett and Satpute, 2013; Buckner, 2012) of the meaning of sensations from the body and world during cognitions, emotions, and perceptions.

It is tempting to assume that the lack of specificity for each emotion category is a function of the coarse spatial and temporal resolution in the resting state brain data (or the brain imaging experiments, for that matter), but even human lesions studies (Feinstein, et al., 2013; Hurlemann, et al., 2009) and studies that electrically stimulate specific neurons in fully conscious humans have not been able to identify specific neural modules for specific emotions [e.g., (Guillory and Bujarski, 2014); for a review of studies, see (Barrett, et al., 2007)]. For instance, consistent with our intrinsic network findings, electrical stimulation of the human brain from intracranial electrodes reveals broadly distributed networks across the cortex, paralimbic and limbic cortex, and subcortex that contribute to the representation of multiple emotional states (Guillory and Bujarski, 2014). A growing evidence from other domains has also failed to find evidence of biologically basic emotions, such as studies of autonomic function in humans (Cacioppo, et al., 2000; Kreibig, 2010), facial expressions in infants and adults (Barrett, 2011; Barrett and Kensinger, 2010; Camras, et al., 2002; Russell, et al., 2003) and studies of vocal acoustics (Bachorowski & Owren, 2003).

A related possibility is that emotions are represented as task-evoked functional brain networks that flexibly combine in a given moment to produce the experience or perception of anger, disgust, fear, happiness, sadness, and so on [as hypothesized (Hamann, 2012)]. Such a hypothesis is not orthogonal to a constructionist interpretation we are offering, although we did not examine task-dependent BOLD data and so did not test this idea explicitly in the present report. For example, if the brain possesses of a set of intrinsic networks that can be understood as performing domain-general operations, then it is possible that each process can be observed as a set of basic processing modes (aka “functional motifs;” (Sporns and Kotter, 2004), arising from the anatomical connections that undergirds each network [aka “structural motifs” (Sporns and Kotter, 2004)]. In this framework, individual instances of anger, disgust, fear, etc, could be understood as high dimensional brain states reflecting neural assemblies within broadly distributed networks, as well as the dynamic interaction of those assemblies (cf. Lindquist and Barrett, 2012; Barrett and Satpute, 2013). Consistent with this prediction, in another study from our lab we used task-evoked functional connectivity and examined the relationship between the intensity of ongoing emotional experiences of anger, sadness, and fear, on the one hand, and the continually fluctuating functional connectivity strength between regions of the salience and default networks on the other (Raz, Touroutoglou et al., under review). Across five samples of subjects, we predicted, and found that the dynamic variation in the functional connectivity between intrinsic networks across time (i.e., changing cohesiveness) constituted a shared mechanism for intense experiences of sadness, fear, and anger. These findings are ultimately consistent with the findings reported herein because they suggest that momentary experiences of emotion are related to the functional coupling of intrinsic connectivity networks.

More broadly, our findings are consistent with an alternative framework for understanding the brain's functional architecture. The fact that peak activations from different emotion categories belong to domain-general intrinsic functional connectivity networks is consistent with a broader *constructionist* view of the mind (Lindquist and Barrett, 2012; Barrett and Satpute, 2013), more generally. A constructionist model of mind-brain correspondence hypothesizes that all mental states emerge (i.e., are "constructed") from the interaction of more basic psychological processes that are not specific to folk psychological distinctions such as "emotion", "cognition", "memory" or "perception". Emotions, cognitive functions and perceptions can be thought of as mental events (prompted by specific experimental tasks, or arising as naturally occurring states) that are constructed from interactions within and between intrinsic networks that compute domain-general functions. A host of neuroscience research findings point towards a constructionist functional architecture of the brain that relies on distributed structure-function mappings. This constructionist approach echoes other debates about modularity throughout neuroscience (e.g., in face perception), which center on whether a phenomenon has dedicated neural modules or is constructed from more domain-general elements (Anderson and Finlay, 2014; Grill-Spector and Weiner, 2014). Taken together, our and other findings stress the need for revisions in the psychological ontologies so that they are consistent with structure and function of the brain [cf. (Anderson, et al., 2013; Barrett and Satpute, 2013; Fox and Friston, 2012; Lindquist and Barrett, 2012; Poldrack, 2010)].

5. Acknowledgement

We thank R. L. Buckner, Harvard University, for providing the data used in the analysis of Sample 1 and Sample 2 and the preprocessing/rs-fcMRI tools. This work was supported by the

National Institutes of Health Director's Pioneer Award (DP1OD003312) to Lisa Feldman Barrett, a National Institute on Aging grant (R01 AG030311-06A1) to Lisa Feldman Barrett and Brad Dickerson, the Shared Instrumentation Grants (1S10RR023401, 1S10RR019307, and 1S10RR023043) from the National Center For Research Resources, and a Harvard University Mind/Brain/Behavior Initiative Postdoctoral Fellowship to Kristen Lindquist. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Center For Research Resources, the National Institutes of Health, or the National Institute on Aging.

6. References

- Anderson, M.L., Finlay, B.L. (2014) Allocating structure to function: the strong links between neuroplasticity and natural selection. *Front Hum Neurosci*, 7:918.
- Anderson, M.L., Kinnison, J., Pessoa, L. (2013) Describing functional diversity of brain regions and brain networks. *Neuroimage*, 73:50-8.
- Andrews-Hanna, J.R., Reidler, J.S., Huang, C., Buckner, R.L. (2010) Evidence for the default network's role in spontaneous cognition. *Journal of neurophysiology*, 104:322-322.
- Barrett, L.F. (2006) Solving the emotion paradox: Categorization and the experience of emotion. *Personality and Social Psychology Review*, 10:20-46.
- Barrett, L.F. (2009) The Future of Psychology: Connecting Mind to Brain. *Perspect Psychol Sci*, 4:326-339.
- Barrett, L.F. (2011) Was Darwin wrong about emotional expressions? *Current Directions in Psychological Science*, 20:400-406.

- Barrett, L.F. (2012) Emotions are real. *Emotion*, 12:413-29.
- Barrett, L.F. (2013) Psychological Construction: The Darwinian Approach to the Science of Emotion. *Emotion Review*, 5:379-389.
- Barrett, L.F. (2014) The conceptual act theory: A précis. *Emotion Review* 6: 292-297.
- Barrett, L.F., Kensinger, E.A. (2010) Context Is Routinely Encoded During Emotion Perception. *Psychological Science*, 21:595-595.
- Barrett, L.F., Lindquist, K.A., Bliss-Moreau, E., Duncan, S., Gendron, M., Mize, J., Brennan, L. (2007) Of mice and men: Natural kinds of emotions in the mammalian brain? A response to Panksepp and Izard. *Perspectives on psychological science: a journal of the Association for Psychological Science*, 2:297-297.
- Barrett, L.F., Satpute, A.B. (2013) Large-scale brain networks in affective and social neuroscience: towards an integrative functional architecture of the brain. *Current Opinion in Neurobiology*, 23:361-372.
- Bickart, K.C., Hollenbeck, M.C., Barrett, L.F., Dickerson, B.C. (2012) Intrinsic amygdala-cortical functional connectivity predicts social network size in humans. *J Neurosci*, 32:14729-41.
- Binder, J.R., Desai, R.H., Graves, W.W., Conant, L.L. (2009) Where is the semantic system? A critical review and meta-analysis of 120 functional neuroimaging studies. *Cerebral cortex*, 19:2767-96.
- Bressler, S.L., Menon, V. (2010) Large-scale brain networks in cognition: emerging methods and principles. *Trends Cogn Sci*, 14:277-90.
- Buckner, R.L. (2012) The serendipitous discovery of the brain's default network. *Neuroimage*, 62:1137-45.

- Buckner, R.L., Andrews-Hanna, J.R., Schacter, D.L. (2008) The brain's default network: anatomy, function, and relevance to disease. *Annals of the New York Academy of Sciences*, 1124:1-38.
- Buckner, R.L., Krienen, F.M., Yeo, B.T. (2013) Opportunities and limitations of intrinsic functional connectivity MRI. *Nat Neurosci*, 16:832-7.
- Bzdok, D., Schilbach, L., Vogeley, K., Schneider, K., Laird, A.R., Langner, R., Eickhoff, S.B. (2012) Parsing the neural correlates of moral cognition: ALE meta-analysis on morality, theory of mind, and empathy. *Brain structure & function*, 217:783-96.
- Cacioppo, J.T., Berntson, G.G., Larsen, J.T., Poehlmann, K.M., Ito, T.A. (2000) The psychophysiology of emotion. . In: Lewis, M., Haviland-Jones, J.M., editors. *Handbook of emotions* 2nd ed. New York: Guilford Press. p pp. 173-191.
- Camras, L.A., Meng, Z., Ujiie, T., Dharamsi, S., Miyake, K., Oster, H., Wang, L., Cruz, J., Murdoch, A., Campos, J. (2002) Observing emotion in infants: facial expression, body behavior, and rater judgments of responses to an expectancy-violating event. *Emotion*, 2:179-93.
- Corbetta, M., Patel, G., Shulman, G.L. (2008) The reorienting system of the human brain: from environment to theory of mind. *Neuron*, 58:306-24.
- Craig, A.D. (2002) How do you feel? Interoception: the sense of the physiological condition of the body. *Nature reviews. Neuroscience*, 3:655-66.
- Decety, J., Jackson, P.L. (2004) The functional architecture of human empathy. *Behavioral and cognitive neuroscience reviews*, 3:71-100.
- Deco, G., Jirsa, V.K., McIntosh, A.R. (2010) Emerging concepts for the dynamical organization of resting-state activity in the brain. *Nature Reviews Neuroscience*, 12:43-56.

- Dickerson, B.C., Eichenbaum, H. (2010) The episodic memory system: neurocircuitry and disorders. *Neuropsychopharmacology*, 35:86-104.
- Ekman, P. (1999) Facial expressions. *Handbook of cognition and emotion*, 16:301-320.
- Feinstein, J.S., Buzzetta, C., Hurlemann, R., Follmer, R.L., Dahdaleh, N.S., Coryell, W.H., Welsh, M.J., Tranel, D., Wemmie, J.A. (2013) Fear and panic in humans with bilateral amygdala damage. *Nature Neuroscience*, 16:270-2.
- Fox, P.T., Friston, K.J. (2012) Distributed processing; distributed functions? *Neuroimage*, 61:407-26.
- Greicius, M.D., Kiviniemi, V., Tervonen, O., Vainionpaa, V., Alahuhta, S., Reiss, A.L., Menon, V. (2008) Persistent default-mode network connectivity during light sedation. *Hum Brain Mapp*, 29:839-47.
- Greicius, M.D., Srivastava, G., Reiss, A.L., Menon, V. (2004) Default-mode network activity distinguishes Alzheimer's disease from healthy aging: evidence from functional MRI. *Proc Natl Acad Sci U S A*, 101:4637-42.
- Grill-Spector, K., Weiner, K.S. (2014) The functional architecture of the ventral temporal cortex and its role in categorization. *Nature reviews. Neuroscience*, 15:536-48.
- Gross, C.T., Canteras, N.S. (2012) The many paths to fear. *Nature reviews. Neuroscience*, 13:651-8.
- Guillory, S.A., Bujarski, K.A. (2014) Exploring emotions using invasive methods: review of 60 years of human intracranial electrophysiology. *Soc Cogn Affect Neurosci*.
- Hamann, S. (2012) Mapping discrete and dimensional emotions onto the brain: controversies and consensus. *Trends Cogn Sci*, 16:458-66.

- Hayes, D.J., Northoff, G. (2011) Identifying a network of brain regions involved in aversion-related processing: a cross-species translational investigation. *Frontiers in integrative neuroscience*, 5:49.
- Hermundstad, A.M., Bassett, D.S., Brown, K.S., Aminoff, E.M., Clewett, D., Freeman, S., Frithsen, A., Johnson, A., Tipper, C.M., Miller, M.B., Grafton, S.T., Carlson, J.M. (2013) Structural foundations of resting-state and task-based functional connectivity in the human brain. *Proc Natl Acad Sci U S A*, 110:6169-74.
- Hurlemann, R., Schlaepfer, T.E., Matusch, A., Reich, H., Shah, N.J., Zilles, K., Maier, W., Bauer, A. (2009) Reduced 5-HT(2A) receptor signaling following selective bilateral amygdala damage. *Soc Cogn Affect Neurosci*, 4:79-84.
- Keltner, D., Ekman, P. (2000) Facial expressions of emotion In: Haviland-Jones, J.M., Lewis, M., editors. *Handbook of Emotions*. New York: Guilford. p 236-249.
- Kober, H., Barrett, L.F., Joseph, J., Bliss-Moreau, E., Lindquist, K., Wager, T.D. (2008) Functional grouping and cortical-subcortical interactions in emotion: a meta-analysis of neuroimaging studies. *NeuroImage*, 42:998-1031.
- Kreibig, S.D. (2010) Autonomic nervous system activity in emotion: A review. *Biological psychology*, 84:394-421.
- Laird, A.R., Fox, P.M., Eickhoff, M., Turner, J.A., Ray, K.L., McKay, D.R., Glahn, D.C., Beckmann, C.F., Smith, S.M., Fox, P.T. (2011) Behavioral interpretations of intrinsic connectivity networks. *Journal of Cognitive Neuroscience*, 23:4022-4037.
- LeDoux, J. (2007) The amygdala. *Curr Biol*, 17:R868-74.
- LeDoux, J. (2012) Rethinking the emotional brain. *Neuron*, 73:653-676.

- Lin, D., Boyle, M.P., Dollar, P., Lee, H., Lein, E.S., Perona, P., Anderson, D.J. (2011) Functional identification of an aggression locus in the mouse hypothalamus. *Nature*, 470:221–226.
- Lindquist, K.A., Barrett, L.F. (2012) A functional architecture of the human brain: emerging insights from the science of emotion. *Trends in Cognitive Sciences*, 16:533-540.
- Lindquist, K.A., Satpute, A., Weber, J., Wager, T.D., Barrett, L.F. (in press) The brain basis of positive and negative emotion: Evidence from a meta-analysis of the human neuroimaging literature. *Cerebral Cortex*
- Lindquist, K.A., Wager, T.D., Kober, H., Bliss-Moreau, E., Barrett, L.F. (2012) The brain basis of emotion: A meta-analytic review. *Behav Brain Sci*, 35:121-43.
- Lohmann, G., Hoehl, S., Brauer, J., Danielmeier, C., Bornkessel-Schlesewsky, I., Bahlmann, J., Turner, R., Friederici, A. (2010) Setting the frame: the human brain activates a basic low-frequency network for language processing. *Cerebral cortex*, 20:1286-92.
- Manoliu, A., Meng, C., Brandl, F., Doll, A., Tahmasian, M., Scherr, M., Schwerthoffer, D., Zimmer, C., Forstl, H., Bauml, J., Riedl, V., Wohlschlager, A.M., Sorg, C. (2013) Insular dysfunction within the salience network is associated with severity of symptoms and aberrant inter-network connectivity in major depressive disorder. *Front Hum Neurosci*, 7:930.
- Mantini, D., Corbetta, M., Romani, G.L., Orban, G.A., Vanduffel, W. (2013) Evolutionarily novel functional networks in the human brain? *J Neurosci*, 33:3259-75.
- Menon, V. (2011) Large-scale brain networks and psychopathology: a unifying triple network model. *Trends Cogn Sci*, 15:483-506.

- Menon, V., Uddin, L.Q. (2010) Saliency, switching, attention and control: a network model of insula function. *Brain structure & function*, 214:655-67.
- Motta, S.C., Goto, M., Gouveia, F.V., Baldo, M.V., Canteras, N.S., Swanson, L. (2009) Dissecting the brain's fear system reveals the hypothalamus is critical for responding in subordinate conspecific intruders. *Proceedings of the National Academy of Sciences*, 106:4870-4875.
- Nelson, S.M., Dosenbach, N.U., Cohen, A.L., Wheeler, M.E., Schlaggar, B.L., Petersen, S.E. (2010) Role of the anterior insula in task-level control and focal attention. *Brain structure & function*, 214:669-80.
- Oosterwijk, S., Lindquist, K.A., Anderson, E.C., Dautoff, R., Moriguchi, Y., Barrett, L.F. (2012) States of mind: Emotions, body feelings, and thoughts share distributed neural networks. *NeuroImage*:2110-2128.
- Panksepp, J. (1998) *Affective Neuroscience: The Foundations of Human and Animal Emotions*. New York. Oxford University Press.
- Pernice, V., Staude, B., Cardanobile, S., Rotter, S. (2011) How structure determines correlations in neuronal networks. *PLoS computational biology*, 7:e1002059.
- Poldrack, R.A. (2010) Mapping mental function to brain structure: How can cognitive neuroimaging succeed? *Perspectives on Psychological Science*, 5:753–761.
- Raichle, M.E. (2010) Two views of brain function. *Trends Cogn Sci*, 14:180-90.
- Rilling, J.K., Barks, S.K., Parr, L.A., Preuss, T.M., Faber, T.L., Pagnoni, G., Bremner, J.D., Votaw, J.R. (2007) A comparison of resting-state brain activity in humans and chimpanzees. *Proceedings of the National Academy of Sciences of the United States of America*, 104:17146-51.

- Russell, J.A., Bachorowski, J.A., Fernandez-Dols, J.M. (2003) Facial and vocal expressions of emotion. *Annual Review of Psychology*, 54:329–349.
- Seeley, W.W., Menon, V., Schatzberg, A.F., Keller, J., Glover, G.H., Kenna, H., Reiss, A.L., Greicius, M.D. (2007) Dissociable intrinsic connectivity networks for salience processing and executive control. *The Journal of neuroscience : the official journal of the Society for Neuroscience*, 27:2349-56.
- Shirer, W.R., Ryali, S., Rykhlevskaia, E., Menon, V., Greicius, M.D. (2012) Decoding Subject-Driven Cognitive States with Whole-Brain Connectivity Patterns. *Cerebral cortex*.
- Smith, S.M., Fox, P.T., Miller, K.L., Glahn, D.C., Fox, P.M., Mackay, C.E., Filippini, N., Watkins, K.E., Toro, R., Laird, A.R., Beckmann, C.F. (2009) Correspondence of the brain's functional architecture during activation and rest. *Proceedings of the National Academy of Sciences of the United States of America*, 106:13040-5.
- Spector, E.R., Smith, S.M., Sibonga, J.D. (2009) Skeletal effects of long-duration head-down bed rest. *Aviat Space Environ Med*, 80:A23-8.
- Sporns, O. (2013) Structure and function of complex brain networks. *Dialogues Clin Neurosci*, 15:247-62.
- Sporns, O., Kotter, R. (2004) Motifs in brain networks. *PLoS Biol*, 2:e369.
- Tomasi, D., Volkow, N.D. (2012) Resting functional connectivity of language networks: characterization and reproducibility. *Mol Psychiatry*, 17:841-54.
- Touroutoglou, A., Bickart, K.C., Barrett, L.F., Dickerson, B.C. (2014) Amygdala task-evoked activity and task-free connectivity independently contribute to feelings of arousal. *Human Brain Mapping*, 35:5316-5327.

- Touroutoglou, A., Hollenbeck, M., Dickerson, B.C., Feldman Barrett, L. (2012) Dissociable large-scale networks anchored in the right anterior insula subserve affective experience and attention. *NeuroImage*, 60:1947-1958.
- Tracy, J.L., Randles, D. (2011) Four Models of Basic Emotions: A Review of Ekman and Cordaro, Izard, Levenson, and Panksepp and Watt. *Emotion Review*, 3:397-405.
- van den Heuvel, M.P., Mandl, R.C., Kahn, R.S., Hulshoff Pol, H.E. (2009) Functionally linked resting-state networks reflect the underlying structural connectivity architecture of the human brain. *Human Brain Mapping*, 30:3127-41.
- van den Heuvel, M.P., Sporns, O. (2013) An anatomical substrate for integration among functional networks in human cortex. *J Neurosci*, 33:14489-500.
- Van Dijk, K.R., Hedden, T., Venkataraman, A., Evans, K.C., Lazar, S.W., Buckner, R.L. (2010) Intrinsic functional connectivity as a tool for human connectomics: theory, properties, and optimization. *Journal of neurophysiology*, 103:297-321.
- Vazdarjanova, A., McGaugh, J.L. (1998) Basolateral amygdala is not critical for cognitive memory of contextual fear conditioning. *Proceedings of the National Academy of Sciences* 95:15003-15007.
- Vincent, J.L., Kahn, I., Snyder, A.Z., Raichle, M.E., Buckner, R.L. (2008) Evidence for a frontoparietal control system revealed by intrinsic functional connectivity. *Journal of neurophysiology*, 100:3328-42.
- Vincent, J.L., Patel, G.H., Fox, M.D., Snyder, A.Z., Baker, J.T., Van Essen, D.C., Zempel, J.M., Snyder, L.H., Corbetta, M., Raichle, M.E. (2007) Intrinsic functional architecture in the anaesthetized monkey brain. *Nature*, 447:83-6.

Vogt, B.A. (2005) Pain and emotion interactions in subregions of the cingulate gyrus. *Nature reviews. Neuroscience*, 6:533-44.

Vytal, K., Hamann, S. (2010) Neuroimaging support for discrete neural correlates of basic emotions: a voxel-based meta-analysis. *Journal of Cognitive Neuroscience*, 22:2864-2885.

Wang, D., Qin, W., Liu, Y., Zhang, Y., Jiang, T., Yu, C. (2014) Altered resting-state network connectivity in congenital blind. *Hum Brain Mapp*, 35:2573-81.


Wilson-Mendenhall, C.D., Barrett, L.F., Barsalou, L.W. (2013) Neural evidence that human emotions share core affective properties. *Psychol Sci*, 24:947-56.

Wilson-Mendenhall, C.D., Barrett, L.F., Simmons, W.K., Barsalou, L.W. (2011) Grounding emotion in situated conceptualization. *Neuropsychologia*, 49:1105-27.


Yeo, B.T., Krienen, F.M., Sepulcre, J., Sabuncu, M.R., Lashkari, D., Hollinshead, M., Roffman, J.L., Smoller, J.W., Zollei, L., Polimeni, J.R., Fischl, B., Liu, H., Buckner, R.L. (2011) The organization of the human cerebral cortex estimated by intrinsic functional connectivity. *J Neurophysiol*, 106:1125-65.

¹For example, Ekman (1999) wrote “It is necessary to posit emotion-specific central nervous system (CNS) activity in my account of basic emotions. The distinctive features of each emotion, including the changes not just in expression but in memories, imagery, expectations and other cognitive activities, could not occur without central nervous system organization and direction. There must be unique physiological [CNS] patterns for each emotion” (p.50). More recently, in


a review of basic emotion theories, Tracy and Randles (2011) wrote that an “agreed-upon gold standard is the presence of neurons dedicated to the emotion’s activation” (p.398).



188x187mm (150 x 150 DPI)



116x75mm (150 x 150 DPI)



171x172mm (150 x 150 DPI)

Figure Legends

Figure 1. The conjunction of discovery maps from meta-analytic activation peaks for happiness, anger, fear, sadness, and disgust at $z(r) = 0.1$ (Sample 1, $N = 89$). In (a), voxels that preferentially correlate with the left superior temporal gyrus (STG) seed are shown in blue, voxels that correlate with left anterior cingulate cortex (ACC) seed are shown in red, and voxels that correlate with the left cerebellum seed are shown in yellow. In (b), voxels that preferentially correlate with left inferior frontal gyrus (IFG) seed are shown in blue, voxels that correlate with right parahippocampal gyrus (PHG) seed are shown in red, and voxels that correlate with the fusiform gyrus seed are shown in yellow. In (c), voxels that preferentially correlate with bilateral amygdala seed regions are shown in blue, voxels that correlate with the right cerebellum seed are shown in red, and voxels that correlate with right insula seed are shown in yellow. In (d), voxels that preferentially correlate with left medial frontal gyrus (medFG) seed are shown in blue, voxels that correlate with the right inferior frontal gyrus (IFG) seed are shown in red, and voxels that correlate with the left caudate seed are shown in yellow. In (e), voxels that preferentially correlate with bilateral inferior frontal gyrus/insula (IFG/insula) seed are shown in blue, voxels that correlate with the left lingual gyrus seed (IFG) are shown in red, and voxels that correlate with the left amygdala seed are shown in yellow. The binarized correlation maps, $z(r) = 0.1$ are overlaid on the 1mm MNI152 T1-standard template image in FSL (<http://www.fmrib.ox.ac.uk/fsl>)

Figure 2. The average strength of intrinsic connectivity, $z(r)$, values between the three seeds associated with each emotion category, the three seeds associated with the default mode network

(DMN) (see supplementary materials), and the control seeds [associated with motor (Mot), auditory (Aud) and vision (Vis) networks; see supplementary materials] in Sample 1 and Sample 2.

Figure 3. A conjunction map for negative emotions. This figure displays a conjunction map of the binarized maps, $z(r) = 0.05$, seeded by the most prominent peak of each negative emotion (i.e., anger; L Inferior Frontal Gyrus seed, sadness; L medial Frontal Gyrus seed, fear; L amygdala seed, and disgust; L Inferior Frontal Gyrus/insula seed).

Table 1. Seed regions of interest

Label	MNI Coordinates		
	x	y	z
Happiness			
Right Superior Temporal Gyrus (R STG)*	48	-55	-4
Left Anterior Cingulate Cortex (L ACC)	-2	43	7
Left Cerebellum	-40	-63	-25
Anger			
Left Inferior Frontal Gyrus (L IFG)*	-45	23	-3
Right Parahippocampal Gyrus (R PHG)	19	-20	-11
Left Fusiform Gyrus	-44	-72	-18
Fear			
Left Amygdala *	-23	-6	-11
Right Amygdala	23	-10	-14
Right Cerebellum	33	-54	-15
Right Insula	43	3	-2
Sadness			
Left medial Frontal Gyrus (L medFG)*	-4	47	32
Right Inferior Frontal Gyrus (R IFG)	40	6	23

Left Caudate head -10 19 -9

Disgust

Right Inferior Frontal Gyrus/Insula 31 5 -4

(rIFG/insula)*

Left Inferior Frontal Gyrus/Insula -26 29 -10

(rIFG/insula)

Left Lingual Gyrus -22 -72 -11

Left amygdala -20 -3 -17

Note. The present paper used a set of three different regions found to be consistently active during each emotion category previously reported in the Vytal and Hamann (2010) meta-analysis of task-induced emotion activations. We selected the three peak activations with the largest cluster of activation for happiness, anger, and sadness. We included four peaks of activation for fear and disgust because the two peaks with the largest cluster of activation were both within the same brain region in different hemispheres. The set of regions for emotions are listed in rank order, based on the size of meta-analytic cluster of activation (*indicates the regions with the largest activation peak for each emotion). All coordinates are referenced to the Montreal Neurological Institute (MNI) coordinate system.

Table 2. Intrinsic connectivity networks of interest used for the goodness of fit analysis

Dorsal and Ventral Salience Networks identified by Tournoutoglou et al.(2012)

1. Dorsal anterior insula network (Dorsal Salience)
2. Ventral anterior insula network (Ventral Salience)

Intrinsic Connectivity Networks identified by Shirer et al.(2012)

1. Insula/dorsal anterior cingulate cortex (Anterior Salience)
 2. Posterior insula (Posterior Salience)
 3. Auditory
 4. Basal ganglia
 5. PCC/Medial prefrontal cortex (dorsal DMN)
 6. Retrosplenial cortex/medial temporal lobe (ventral DMN)
 7. Language
 8. Left dorsolateral prefrontal cortex/Left parietal lobe (Left executive control network, ECN)
 9. Right dorsolateral prefrontal cortex /Right Parietal Lobe (Right executive control network, ECN)
 10. Intraparietal sulcus/Frontal Eye Field (Visuospatial)
 11. Precuneus
 12. Primary visual cortex,V1
 13. Secondary visual cortex,V2 (High-level visual)
 14. Sensorimotor
-

Table 3. The goodness-of-fit values for the emotion discovery maps

Reference Networks	Happiness		Anger		Fear		Sadness		Disgust	
	R STG	L ACC	L IFG	R PHG	L Amygdala	R Amygdala	R Cerebellum	R Insula	L medFG	R IFG/Insula
Dorsal Salience							0.28		0.11	0.10
Ventral Salience	0.35		0.11					0.18		
Anterior Salience			0.12				0.16			
Posterior Salience							0.22			
Dorsal DMN	0.40							0.33	0.14	
Basal Ganglia				0.11	0.12				0.19	0.11
Language			0.21							
L Executive Control							0.21			
Visuospatial	0.16							0.17		
Auditory							0.32			
Primary Visual						0.12				0.31
High-level Visual	0.11			0.24		0.29				0.36

Note: goodness-of-fit values lower than 0.1 are not shown in the table for ease of viewing